arXiv:hep-th/9305034v1 10 May 1993

UT-642
February 1, 2008

Lattice Topological Field Theory and First Order Phase Transition[*]

Naoki Sasakura[⋆][†]

Department of Physics, University of Tokyo, Tokyo, 113, Japan

ABSTRACT

Carrying out perturbations around a lattice topological field theory in two dimensions, we show that it is on a first order phase transition fixed point with multiplicity $n(n-1)/2$, where n is the number of its independent physical observables. We discuss about the order parameters and the finite size effect for the free energy. The finite size effect is described by the topological field theory. We investigate also the renormalization group flow near the fixed point, and show that the flow agrees with that of the Nienhuis-Nauenberg criterion.

---

[*] Work supported by the Grant-in-Aid for Scientific Research from the Ministry of Education No. 04-1324.

[⋆] E-mail address: SASAKURA@TKYUX.PHYS.S.U-TOKYO.AC.JP

[†] A Fellow of the Japanese Society for the Promotion of Science for Japanese Junior Scientists.

It is a well known fact that a continuum field thoery can be defined on a lattice along a renormalization group flow where the limit is a fixed point on a second order phase transition surface. The correlation length diverges on the fixed point. On the other hand there is another possibility for a fixed point of a renormalization group flow, where the correlation length vanishes. Are there any field theories corresponding to such a zero-correlation-length fixed point? Since the correlation length vanishes on such a fixed point, we expect that the correlation functions of some local observables are independent of their positions. This would be a characteristic nature of topological field theories. Recently some lattice topological field theories have been constructed[1−14]. The specific nature of these lattice theories is that their physical observables are completely independent of the lattice structures on which they are defined. In this letter we investigate how the system is influenced by small changes of the local weights of the lattice topological field theories in two dimensions[1,4,13,14], and investigate what kind of points they reside on.

The basic elements of the two-dimensional topological lattice theory are the tensor C^T_{ijk} and the metric g^{ij} of complex numbers satisfying the following constraints[14]:

$$\begin{aligned} C^T_{ijp} g^{pp'} C^T_{p'kl} &= C^T_{jkp} g^{pp'} C^T_{p'li}, \\ g^{ij} &= g^{ii'} g^{jj'} C^T_{i'kl} g^{kk'} g^{ll'} C^T_{j'l'k'}, \end{aligned} \tag{1}$$

where the Roman indices run from 1 to M, and the tensor and the metric are invariant under the cyclic permutations of the indices. Consider a triangulated two-dimensional oriented surface with genus g. We assign C^T_{ijk} to each triangle, where i, j, k represent the degrees of freedom on its three edges. And we assign g^{ij} to each edge and connect the tensors on the two triangles which have the common edge. Then we will obtain a numerical value for the triangulated surface and call it the partition function[1,4,13,14] of the surface:

$$Z^T_g \equiv C^T_{i_1 i_2 i_3} g^{i_3 i_4} \cdots . \tag{2}$$

The condition (1) guarantees the invariance of the partition function (2) under the

Mateev moves of the triangulation[14], and hence it is independent of the triangulation of the surface[1,4,13,14].

We make a small shift of the tensor δC_{ijk} from the original value corresponding to the topological theory:

$$C_{ijk} = C^T_{ijk} + \delta C_{ijk}.$$

We define the partition function by substituting C^T_{ijk} with C_{ijk} in (2). Since the tensors do not necessarily satisfy the condition (1) anymore, the partition function depends on the triangulation. Thus from now on we consider certain triangulations of the surface. We will calculate the partition function perturbatively in δC_{ijk}. The first order terms are given by

$$Z^{(1)}_g = \sum_x \delta C_{ijk} \langle P^{ijk}(x) \rangle_g, \tag{3}$$

where the operator $P^{ijk}(x)$ denotes the puncture operator[14] of the triangle at x with indices i, j, k on its edges, and the one-point function $\langle P^{ijk}(x) \rangle_g$ is that of the original lattice topological field theory on the surface with genus g. Because of the topological nature, the one point function $\langle P^{ijk}(x) \rangle_g$ is independent of the position x, and hence the summation over the position x can be trivially done. Moreover all the puncture operators P^{ijk} are not linearly independent in the correlation functions, and they are equal to certain linear combinations of the physical puncture operators P^i $(i = 1, \cdots, n)$ $(n \leq M)$[14]. Thus the equation (3) is expressed as

$$\begin{aligned} Z^{(1)}_g &= N C_l \langle P^l \rangle_g, \\ \delta C_l &\equiv \delta C_{ijk} A^{ijk}_l, \end{aligned} \tag{4}$$

where N is the total number of the triangles on the triangulated surface, and A^{ijk}_l are the projections to the physical puncture operators. We did not change the metric tensor g^{ij}. This is because the change in the metric tensor will change only the definition of δC_i in (4) and it will not affect the discussions below.

It is not convenient to take the thermo-dynamical limit $N \to \infty$ of the perturbed partition function $Z_g^T + Z_g^{(1)}$, as can be seen from the higher orders. The p-th order is a summation over all the selections of p triangles among the N triangles on the surface. The leading term in N comes from the cases where any of the p triangles are not next neighbors. Since the correlation functions are independent of their positions if they are not next neighbors, we obtain

$$Z_g^{(p)} = \frac{N^p}{p!}\langle(\delta C_l P^l)^p\rangle + O(N^{p-1}(\delta C)^p).$$

Thus the partition function is obtained in a compact way as

$$\begin{aligned} Z_g &= Z_g^T + \sum_{p=1}^{\infty} Z_g^{(p)} \\ &= \langle \exp(N\delta C_i P^i)\rangle_g + (\text{lower orders}). \end{aligned} \tag{5}$$

One can show iteratively that the partition function can be generally expressed in the form $\langle\exp(W)\rangle_g$ in a perturbative treatment, and that $N\delta C_i P^i$ in (5) is the first order of a cluster expansion of W. The p-th order of the cluster expansion comes from p neighboring triangles. Thus the terms higher than the first one depend generally on the lattice structure, and hence they are not universal. In this paper we assume that W can be treated perturbatively in δC_{ijk}.

The calculation rules in the two-dimensional lattice topological field theories are very simple in the standard basis[14]. The physical puncture operators satisfy the following rules :

$$\begin{aligned} P^i P^j &\sim \delta^{ij} P^i \quad \text{(Fusion Rule)}, \\ v^i &\equiv \langle P^i\rangle_0, \\ H &= \sum_{i=1}^{n} \frac{P^i}{v^i}, \\ \langle \mathcal{O}\rangle_g &= \langle \mathcal{O} H^g\rangle_0, \end{aligned}$$

where $\langle \mathcal{O}\rangle_g$ denotes the correlation function of operator $\mathcal{O}$ on a surface of genus g. And the v^i's take only discrete vales, i.e., squares of integers[13]. Using the above rules, the numerical value of the partition function (5) is

$$\begin{aligned} Z_g &= \sum_{p=0}^{\infty} Z_g^{(p)} \\ &= \sum_{i=1}^{n} \langle P^i \rangle_g \exp(N\delta C_i), \\ \langle P^i \rangle_g &= (v^i)^{1-g}. \end{aligned} \tag{6}$$

Thus in the thermo-dynamical limit $N \to \infty$ the free energy per triangle is

$$\begin{aligned} f &= - \lim_{N\to\infty} \frac{1}{N} \log(Z_g) \\ &= - \max_i \delta C_i \end{aligned} .$$

This result shows that the position of the lattice topological field theory ($\delta C_i = 0$) is a multi-critical first order phase transition point with multiplicity $n(n-1)/2$, around which there are n different phases.

The order parameters of the phase transitions are given by the one-point functions of the puncture operators. Define the one-point functions as follows:

$$\langle P^l \rangle_g^{\delta C} \equiv \frac{A^l_{ijk} Z(P^{ijk})_g}{Z_g},$$

where $Z(P^{ijk})_g$ is the partition function of the triangulated surface with a boundary of a triangle with indices i, j, k on its edges. In the same way as the above calculations for the partition function, one obtains in the first order of the cluster expansion

$$\langle P^l \rangle_g^{\delta C} = \frac{\sum_{i=1}^{n} (\delta^{li} + O(\delta C)^{li}) \langle P^i \rangle_g \exp(N\delta C_i)}{\sum_{i=1}^{n} \langle P^i \rangle_g \exp(N\delta C_i)}.$$

Here the $O(\delta C)^{li}$ terms are the first order terms of the cluster expansion for the puncture operator, which depend on the local lattice structure. If we are in the

m-th phase, i.e., δC_m is larger than the others, we obtain in the thermo-dynamical limit

$$\langle P^l \rangle_g^{\delta C} = \delta^{ml} + O(\delta C)^{ml},$$

which shows they are good order parameters.

As can be seen in the above calculations, the one-point functions of the puncture operators of the lattice topological field theory do not appear anywhere after taking the thermo-dynamical limit. But they appear as the finite size effect, and in the m-th phase, the asymptotic form of the free energy per triangle is obtained from (6) as

$$f \sim -\delta C_m - \frac{1}{N} \log \langle P^m \rangle_g.$$

One would notice that the coefficient and the topology dependence of the finite size effect is calculable by the topological field theory, and that the $1/N$ finite size effect appears only when $g \neq 1$.

Next we will discuss the renormalization group flow near the multi-critical point. A partition function must be invariant under a renormalization. Suppose that the number of triangles N change to N' under one renormalization step. The most natural way to realize the invariance of the partition function (5) under the renormalization step is to define the renormalized coupling $\delta C'_i$ such that $N \delta C_i = N' \delta C'_i$. This renormalization group flow agrees with that of the Nienhuis-Nauenberg criterion[15], which gives the renormalization group flow near a discontinuous fixed point, and shows that the multi-critical point is the ultraviolet fixed point of the flow.

Simple examples of such topological fixed points[13] can be found in the parameter space of the ferromagnetic Potts model. Consider Potts spins $i, j, \cdots \in \{1, \cdots, Q\}$ located on the sites of a triangular lattice and interacting among nearest neighbors. This model can be defined by a metric and a rank-three tensor as follows:

$$\begin{aligned} g^{(ij)(kl)} &= \delta^{il}\delta^{jk}W(i,j), \\ C_{(ij)(kl)(mn)} &= \kappa\delta_{jk}\delta_{lm}\delta_{ni}W(i,j)W(k,l)W(m,n), \\ W(i,j) &= \exp\left(\frac{\delta_{ij}-1}{3T}\right), \end{aligned}$$

where κ is a constant and T is the temperature. Those tensors satisfy the topological condition (1) at $T=0$ and $T=\infty$ by choosing κ appropriately. At $T=0$ the model is a lattice topological field theory of $n=Q$ and $v^i=1$ $(i=1,\cdots,Q)$. At $T=\infty$ the model is of $n=1$ and $v^1=Q^2$, and the partition function depends on the topology of the surface. At the both points the correlation length is zero, and the lattice topological field theories are realized on the zero-correlation-length fixed points.

In this letter we clarified some universal natures of the topological fixed points by the perturbative treatment. A fixed point on a first order phase transition surface is a zero-correlation-length fixed point. In fact we have shown that a lattice topological field theory with more than one physical observable is on a multi-critical first order phase transition point. Any point near a first order phase transition surface would be transformed to the neighborhood of a fixed point by a renormalization group transformation. From this view point we believe that this letter suggests the possibility that lattice topological field theories would be useful to understand and classify the first order phase transitions. Thus it would be an interesting challenge to investigate whether any lattice topological field theories can also describe the non-trivial first order phase transition points of the Potts models with $Q>4$.

Acknowledgements: We would like to thank H. Kawai, T. Yukawa, Y. Kitazawa, K.I. Izawa and N. Kawamoto for valuable discussions, comments, criticism and encouragement. We would like to thank also the members of the elementary particle sections of Tokyo University and Kyoto University for their hospitality. And we would like to thank H. Kawai for reading this manuscript as well as N. Kawamoto for checking the manuscript.

REFERENCES

1. J.F. Wheater, *Phys. Lett.* **B223** (1989), 451.
2. R. Dijkgraaf and E. Witten, *Commun. Math. Phys.* **129** (1990), 393.
3. Al.R. Kavalov and R.L. Mkatchyan, *Phys. Lett.* **242B** (1990), 429.
4. T. Jònsson, *Phys. Lett.* **B265** (1991), 141.
5. V.G. Turaev and O.Y. Viro, "*State Sum Invariant of 3-Manifolds and Quantum 6j-Symbols*," preprint (1990).
6. H. Ooguri and N. Sasakura, *Mod. Phys. Lett* **A6** (1991), 3591.
7. S. Mizoguchi and T. Tada, *Phys. Rev. Lett.* **68** (1992), 1795.
8. F. Archer and R.M. Williams, *Phys. Lett.* **B273** (1991), 438.
9. B. Durhuus, H. Jacobsen and R. Nest, *Nucl. Phys.(Proc. Suppl.)* **25A** (1992), 109.
10. D.V. Boulatov, *Mod. Phys. Lett* **A7** (1992), 1629.
11. H. Ooguri, *Mod. Phys. Lett* **A7** (1992), 2799.
12. F.J. Archer, *Phys. Lett.* **B295** (1992), 199.
13. C. Bachas and P.M.S. Petropoulos, *Commun. Math. Phys.* **152** (1993), 191.
14. M. Fukuma, S. Hosono and H. Kawai, preprint CLNS 92/1173.
15. B. Nienhuis and M. Nauenberg, *Phys. Rev. Lett.* **35** (1975), 477.